

VACCINATION SCHEDULE

Sprache: Englisch

Vaccinations should be administered at the earliest possible age.

An assessment of vaccination status is recommended at all ages. Any missed vaccinations should be administered immediately, in accordance with the recommendations for the relevant age group.

German Standing Committee on Vaccination (STIKO), recommendations, 2019/20. www.stiko.de

ROBERT KOCH INSTITUT



VACCINATION	AGE	(in weeks)			(in months)		(in years)			Adolescents		Adults			
		6	2	3	4	11-14	15-23	2-4	5-6	7-8	9-14	15-16	17	from 18	from 60
		U4			U6	U7	U7/8	U9	U10	J1					
Rotaviruses		G1 ^a	G2	(G3)											
Tetanus			G1	G2	G3	G4	N		A1	N	A2	N	A (N if required) ^f		
Diphtheria			G1	G2	G3	G4	N		A1	N	A2	N	A (N if required) ^f		
Whooping cough Pertussis			G1	G2	G3	G4	N		A1	N	A2	N	A3 ^f	N if required	
Hib <i>Haemophilus influenzae</i> Type b			G1	G2 ^b	G3	G4	N								
Polio Poliomyelitis			G1	G2 ^b	G3	G4	N			A1	N	N if required			
Hepatitis B			G1	G2 ^b	G3	G4	N								
Pneumococcal disease ^c			G1		G2	G3	N							S ^g	
Meningococcal disease C						G1 (from 12 months)		N							
Measles						G1	G2	N					S ^e		
Mumps Rubella						G1	G2	N							
Chicken pox Varicella						G1	G2	N							
HPV Human papillomaviruses								G1 ^d G2 ^d		N ^d					
Herpes zoster												G1 ^h	G2 ^h		
Flu Influenza												S (annually)			

Dates of next vaccinations



EXPLANATORY NOTES

- G** PRIMARY IMMUNISATION (up to 4 doses of vaccine G1 – G4)
- S** STANDARD VACCINATION
- A** BOOSTER VACCINATION
- N** CATCH-UP VACCINATION (primary immunisation of persons not yet vaccinated or completion of an incomplete vaccination series)
- U** Well-child visit
- J** Well-adolescent visit (J1 aged 12–14 years)

- a** The 1st vaccine dose should be administered from the age of 6 weeks. Depending on the type of vaccine, 2 or 3 doses are required at intervals of a minimum of 4 weeks.
- b** This dose is not required if a monovalent vaccine is administered.
- c** Premature infants receive an additional dose of vaccine at the age of three months, i.e. a total of 4 doses.
- d** Standard vaccination for girls and boys aged 9 to 14. For catch-up vaccinations and number of vaccine doses see the summary of product characteristics.
- e** Single dose vaccination for all individuals ≥ 18 years with unclear vaccination status and who were born after 1970, who have not been vaccinated or only received one vaccination as a child. Preferably with an MMR vaccine.

- f** Td booster vaccination every 10 years. The next Td vaccination that is due is to be administered as a single dose vaccination in the form of Tdap or, if indicated, in the form of a Tdap-IPV combination vaccination.
- g** Vaccination with polysaccharide vaccine, booster vaccination only recommended for specific indications.
- h** Two vaccinations with the adjuvanted herpes zoster vaccine administered at least 2 but no more than 6 months apart.